

Niacin Metabolism and Parkinson's Disease

Tetsuhito FUKUSHIMA¹

¹Department of Hygiene & Preventive Medicine, Fukushima Medical University School of Medicine, Fukushima, Japan

Abstract

Epidemiological surveys suggest an important role for niacin in the causes of Parkinson's disease, in that niacin deficiency, the nutritional condition that causes pellagra, appears to protect against Parkinson's disease. Absorbed niacin is used in the synthesis of nicotinamide adenine dinucleotide (NAD) in the body, and in the metabolic process NAD releases nicotinamide by poly(ADP-ribosyl)ation, the activation of which has been reported to mediate 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced Parkinson's disease. Recently nicotinamide N-methyltransferase (EC2.1.1.1) activity has been discovered in the human brain, and the released nicotinamide may be methylated to 1-methylnicotinamide (MNA), via this enzyme, in the brain. A deficiency in mitochondrial NADH:ubiquinone oxidoreductase (complex I) activity is believed to be a critical factor in the development of Parkinson's disease. MNA has been found to destroy several subunits of cerebral complex I, leading to the suggestion that MNA is concerned in the pathogenesis of Parkinson's disease. Based on these findings, it is hypothesized that niacin is a causal substance in the development of Parkinson's disease through the following processes: NAD produced from niacin releases nicotinamide via poly(ADP-ribosyl)ation, activated by the hydroxyl radical. Released excess nicotinamide is methylated to MNA in the cytoplasm, and superoxides formed by MNA via complex I destroy complex I subunits directly, or indirectly via mitochondrial DNA damage. Hereditary or environmental factors may cause acceleration of this cycle, resulting in neuronal death.

Key words: nicotinamide N-methyltransferase, 1-methylnicotinamide, poly(ADP-ribosyl)ation, mitochondria, complex I

Introduction

Because prokaryotes proliferate by cell division, aging of the individual cell does not occur. In eucaryotes, however, fatal aging occurs. Aging of the neuron begins immediately after its differentiation. The presence of mitochondria, especially their electron transport systems, makes possible an enormous energy supply in eucaryotic cells, and the coenzyme, nicotinamide adenine dinucleotide (NAD), may play an important role in their functioning (1). Contrarily, is it too reckless to attribute the cause of the fatal aging which occurs in eucaryotic cell to the existence of mitochondria and NAD?

In advanced countries, the prevalence of Parkinson's disease is so high that it is classified as one of the diseases of

the aged. Any causal substance for Parkinson's disease will therefore be, not a special compound, but a common one, such as a food nutrient. A number of researchers have reported a relationship between niacin intake and Parkinson's disease. Some findings have indicated that niacin is neuroprotective (2, 3), whereas others have not supported it and niacin showed no clear relation with Parkinson's disease (4–6). NADH has been reported to relieve some of the symptoms of Parkinson's disease, but it did not influence the tissue content of dopamine in vivo experiment (7). Because in advanced countries, enough amounts of niacin are already commonly consumed, these case-control or follow-up studies might not offer any meaningful results regarding the relation between Parkinson's disease and niacin.

Neurotoxicity of niacin has not been reported by the epidemiological study yet, but other several epidemiological reports have indirectly indicated a contributory role for niacin in the pathogenesis of Parkinson's disease. Parkinson's disease is very low in Africa and China (8) where pellagra, caused by niacin deficiency, is a common complaint. Isoniazid, an anti-tubercular agent, relieves the symptoms of Parkinson's disease (9), but induces pellagra as a side effect. Parkinson's disease patients

Received May 5, 2004/Accepted Sep. 24, 2004
Reprint requests to: Tetsuhito FUKUSHIMA
Department of Hygiene & Preventive Medicine, Fukushima Medical University School of Medicine, 1 Hikarigaoka, Fukushima 960-1295, Japan
TEL: +81(24)-547-1173, FAX: +81(24)-547-1174
E-mail: t-fuku@fmu.ac.jp

consume less than average alcohol (10), while heavy drinking is associated with pellagra. Consequently, the people who tend to suffer from niacin deficiency might tend not to suffer from Parkinson's disease. Because, in advanced countries, sufficient niacin to prevent pellagra is commonly consumed, my knowledge of niacin metabolism in the living body may provide evidence consistent with this hypothesis. In this paper the possible role of niacin in the pathogenesis of Parkinson's disease is investigated through a detailed discussion of niacin metabolism.

1. Background of the hypothesis: niacin, mitochondrial dysfunction and Parkinson's disease

Genetic inheritance does not play a significant role in idiopathic Parkinson's disease of advanced age (11) or may play a part of the pathogenesis suggested by the discoveries of the causative genes of familial Parkinson's disease, such as alpha-synuclein, parkin and DJ-1 (12). In the meantime, environmental chemical compounds have been suspected of causing Parkinson's disease since 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) was discovered to cause Parkinson's disease artificially (13). It has been reported that the urinary excretion of 1-methylnicotinamide (MNA) is considerably raised and the excretion of its catabolic product, pyridone, is reduced in Parkinson's disease patients after nicotinamide administration (14). Nicotinamide methylation in the bodies of Parkinson's disease patients is accelerated. Human neurons express transporters, such as hOCT2, that translocate MNA from interstice to neuron (15). MNA or nicotinamide could be transported from blood to brain by specific transport mechanisms (16, 17), which may effect a specific phenomenon in the neuron.

Mitochondrial dysfunction has been implicated in the pathogenesis of several neurodegenerative diseases, including Parkinson's disease (18, 19), and the accumulation of deletions and point mutations in the mitochondrial genome has been a focus of attention (20, 21). Several researchers have reported that a deficiency of NADH: ubiquinone oxidoreductase (complex I) in the mitochondria plays a role in the pathogenesis of Parkinson's disease (22–24). The 30-, 25- and 24-kDa subunits of complex I were decreased in the striata of patients who died of Parkinson's disease (25). More than half of mitochondrially encoded polypeptides form part of the complex I, and mitochondrial polymorphisms are expected to reduce the risk of Parkinson's disease (26). Several researchers have measured respiratory enzyme activities or mitochondrial DNA (mDNA) in the muscle or platelets of Parkinson's disease patients to evaluate whether this change is specific to dopaminergic neurons or is the phenotypically relevant consequence of a widespread failure of the mitochondrial oxidative phosphorylation system (27, 28). They found no difference between Parkinson's disease and control groups. The acceleration of complex I subunit deficiency should therefore occur only in the brain of Parkinson's disease patients, and this specific mechanism of the neurons aroused my interest.

2. Metabolism of NAD with poly(ADP-ribosylation)

Most nicotinic acid exists in animals and plants in the form of nicotinamide, but it is absorbed in the intestines in the form of nicotinic acid, following deamination. It is used in the synthesis of NAD in the human body, and subsequently, in the metabolic process, NAD releases nicotinamide by poly(ADP-ribosylation). This is methylated in the liver and is excreted into the urine as MNA.

DNA damage activates a nuclear enzyme poly(ADP-ribose) synthetase that facilitates DNA repair, and this enzyme activity can provide an early index of DNA damage following neurotoxic insults (29). Excessive activation of this enzyme can, however, deplete tissue stores of NAD, leading to cell death with the depletion of ATP (30). Poly(ADP-ribose) synthetase inhibitors and poly(ADP-ribose) synthetase gene deletion induced dramatic neuroprotection in experimental animals (31). Nitric oxide (via hydroxyl radical) stimulates auto-ADP-ribosylation of glyceraldehyde-3-phosphate dehydrogenase (32–34). Several researchers have attempted to attenuate free radical mediated cerebral damage by inhibition of poly(ADP-ribose) synthetase (35, 36) or by supplementation of niacin (37). Poly(ADP-ribose) synthetase activation mediates MPTP neurotoxicity (38), and its inhibitors protect against MPTP-induced depletion of striatal dopamine (39) or brain NAD and ATP (40). But this is to pay attention to only one consequence of ADP-ribosylation. Poly(ADP-ribosylation) also results in the release from NAD of nicotinamide, which is methylated to MNA in the body. My interest has been in this other consequence of ADP-ribosylation.

3. Nicotinamide methylation in the brain

The occurrence of nicotinamide N-methyltransferase (NAMT) activity in the mammalian brain has been disputed. Shibata (41) reported NAMT activity in the rat brain and liver of 0 (not detected) and 0.282 nmol/mg·h, respectively, and Sano et al. (42), of 0 (not detected) and 5.19 nmol/mg·h, respectively. Both used high performance liquid chromatography with fluorescence detection. Seifert et al. (43), however, reported NAMT activity in the rat brain and liver of 0.4 and 18.6 nmol/mg·h, respectively, using thin-layer chromatography with radioisotope. Early attempts to detect NAMT activity used thin-layer chromatography with radioisotope or high performance liquid chromatography with fluorescence detection. Specificity and sensitivity using these earlier methods were much lower than with our method of gas chromatographic-mass spectrometric analysis in a selected ion monitoring mode. Using gas chromatographic-mass spectrometric analysis in a selected ion monitoring system, the NAMT activities of rat brain and liver were assayed at 0.30 nmol/mg·h and 0.51 nmol/mg·h, respectively (44).

Recently several researchers have reported about the NAMT protein in human brain in relation to the etiology of Parkinson's disease. The relative level of NAMT protein in the lumbar cerebrospinal fluid in Parkinson's disease patients was higher than that in control (45). Expression of NAMT in the parkinsonian cerebella was higher than in the control group (46). From these results the excess NAMT in the central

nervous system might be implicated in the Parkinson's disease pathogenesis.

4. Neurotoxicity of MNA via damage to complex I

Neurotoxicity of MNA was reported using cultured neuroblastoma cells (47), but was not followed up, apart from our research investigating the mechanism of neurotoxicity of MNA. MNA injection in the rat substantia nigra pars compacta significantly decreased dopamine content in the striatum (44). MNA is able to destroy several of the subunits of cerebral complex I, especially 30-kDa protein (48). This subunit is a transmembranous (mitochondrial inner membrane) and iron-sulfur protein which is sensitive to lipid peroxidation (49). NADH oxidation and lipid peroxidation by MNA via rat brain submitochondrial particles (SMP), under conditions of pH ranging from pH 6.0 to 10.0, were verified. The pH optimum for NADH oxidation was 9.0 and the pH optimum for peroxidation by MNA of the lipid composing SMP was also 9.0. The lipid peroxidation in this assay was suppressed by superoxide dismutase (44). Using the paraquat model, it was demonstrated that the lipid peroxidation was caused by the hydroxyl radical which was formed from the superoxide radical (48).

Thus the superoxide anion formed by NADH:MNA oxidation-reduction reaction via complex I might be involved in the etiology of Parkinson's disease by means of direct destruction of complex I subunits, or by an indirect effect via mDNA destruction. Because the human complex I consists of at least 36 nuclear- and 7 mitochondrial-encoded subunits (50, 51), the effect of superoxides formed by MNA on mDNA mutations requires further examination and clarification.

5. Conclusions

NAMT activity in the brain may convert nicotinamide to MNA and by this means damage the nigro-neostriatal dopaminergic neurons. The superoxide anion formed by MNA via the mitochondria may be involved in the etiology of Parkinson's disease. Fig. 1 demonstrates my hypothesis of the process of MNA neurotoxicity. NAD produced from niacin releases nicotinamide via poly(ADP-ribose)ylation, activated by the hydroxyl radical. Released excess nicotinamide is methylated to MNA in the cytoplasm, and superoxides formed by MNA via complex I destroy complex I subunits directly, or indirectly via mDNA damage. Hereditary or environmental factors may cause acceleration of this cycle, resulting in neuronal death. At the level of niacin deficiency there are few nicotinamide methylated to MNA in the living body. But now in advanced countries, enough amounts of niacin are already commonly consumed, and its intake by most people might be over the threshold value causing Parkinson's disease. Aging and hereditary factors should affect individual responses to Parkinson's disease.

6. Perspectives

This hypothesis may have important implications for Parkinson's disease prevention.

Because maize (*Zea mays*) contains niacytin, which humans cannot utilize as niacin, and because maize is also low in tryptophan but abundant in leucine, which inhibits quinolinate phosphoribosyl transferase, the key enzyme for converting from tryptophan to NAD, niacin deficiency is observed in peoples whose main energy source is maize (52). When various countries in the world are sorted hierarchically by their main energy source in food consumption, South Africa, Zimbabwe and




Fig. 1 Hypothesized process of MNA neurotoxicity. MNA, 1-methylnicotinamide; mDNA, mitochondrial DNA; NAD, nicotinamide adenine dinucleotide.

Japan are classified as a maize group (53). The prevalence or mortality rate of Parkinson's disease in these countries, South Africa (54), Zimbabwe (55) and Japan (56), is relatively low. In China, Keshan disease is prevalent in the maize production districts where selenium intake is very low. Niacin deficiency is also found in Keshan disease prevalent districts. Correlation coefficients for the prevalence of Parkinson's disease, maize yield, niacin intake, and selenium intake by province in China were analyzed (57), and a positive correlation was observed between selenium intake and niacin intake. A negative association was found between maize production and niacin intake and between maize production and prevalence of Parkinson's disease. Further, a retrospective study in Japan showed a preventive effect of maize on mortality from Parkinson's disease (58). These results suggest that maize, as a substitute for

other grain crops, can be expected to prevent Parkinson's disease. Of course, the balance of nutrients is important to prevent pellagra.

Today, vitamins are ingested in excess in advanced countries, including Japan (59), while evidence for the long-term safety of their excessive intake is weak. In the light of the above argument, it may be necessary to revisit the recommended amount of niacin for daily consumption.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research (14657090) from the Japanese Ministry of Education, Culture, Sports, Science and Technology.

References

- (1) Scheffler IE. Mitochondria make a come back. *Adv Drug Deliv Rev.* 2001; 49: 3–26.
- (2) Fall PA, Fredrikson M, Axelson O, Granerus AK. Nutritional and occupational factors influencing the risk of Parkinson's disease: a case-control study in southeastern Sweden. *Mov Disord.* 1999; 14: 28–37.
- (3) Hellenbrand W, Boeing H, Robra BP, Seidler A, Vieregge P, Nischan P, Joerg J, Oertel WH, Schneider E, Ulm G. Diet and Parkinson's disease. II: A possible role for the past intake of specific nutrients. Results from a self-administered food-frequency questionnaire in a case-control study. *Neurology.* 1996; 47: 644–650.
- (4) Abbott RD, Ross GW, White LR, Sanderson WT, Burchfiel CM, Kashon M, Sharp DS, Masaki KH, Curb JD, Petrovitch H. Environmental, life-style, and physical precursors of clinical Parkinson's disease: recent findings from the Honolulu-Asia Aging Study. *J Neurol.* 2003; 250 (Suppl 3): III30–39.
- (5) Johnson CC, Gorell JM, Rybicki BA, Sanders K, Peterson EL. Adult nutrient intake as a risk factor for Parkinson's disease. *Int J Epidemiol.* 1999; 28: 1102–1109.
- (6) Ross GW, Abbott RD, Petrovitch H, Morens DM, Grandinetti A, Tung KH, Tanner CM, Masaki KH, Blanchette PL, Curb JD, Popper JS, White LR. Association of coffee and caffeine intake with the risk of Parkinson disease. *JAMA.* 2000; 283: 2674–2679.
- (7) Pearl SM, Antion MD, Stanwood GD, Jaumotte JD, Kapatos G, Zigmund MJ. Effects of NADH on dopamine release in rat striatum. *Synapse.* 2000; 36: 95–101.
- (8) Tanner CM. The role of environmental toxins in the etiology of Parkinson's disease. *Trends Neurosci.* 1989; 12: 49–54.
- (9) Gershanik OS, Luquin MR, Scipioni O, Obeso JA. Isoniazid therapy in Parkinson's disease. *Mov Disord.* 1988; 3: 133–139.
- (10) Fujii C, Harada S, Ohkoshi N, Hayashi A, Yoshizawa K. Study on Parkinson's disease and alcohol drinking. *Nihon Arukoru Yakubutsu Igakkai Zasshi.* 1998; 33: 683–691.
- (11) Tanner CM, Ottman R, Goldman SM, Ellenberg J, Chan P, Mayeux R, Langston JW. Parkinson disease in twins: an etiologic study. *JAMA.* 1999; 281: 341–346.
- (12) Baptista MJ, Cookson MR, Miller DW. Parkin and alpha-synuclein: opponent actions in the pathogenesis of Parkinson's disease. *Neuroscientist.* 2004; 10: 63–72.
- (13) Piccinin GL, Piccirilli M, Finali G, Stefano E. MPTP: a new chapter in the history of Parkinson's disease. *Riv Neurol.* 1989; 59: 103–107.
- (14) Williams AC, Pall HS, Steventon GB, Green S, Buttum S, Molloy H, Waring RH. N-methylation of pyridines and Parkinson's disease. In: Narabayashi H, Nagatsu T, Yanagisawa N, Mizuno Y, editors. *Advances in Neurology*, vol. 60. New York: Raven Press, 1993: p. 194–196.
- (15) Busch AE, Karbach U, Miska D, Gorboulev V, Akhoundova A, Volk C, Arndt P, Ulzheimer JC, Sonders MS, Baumann C, Waldegger S, Lang F, Koepsell H. Human neurons express the polyspecific cation transporter hOCT2, which translocates monoamine neurotransmitters, amantadine, and memantine. *Mol Pharmacol.* 1998; 54: 342–352.
- (16) Köppen A, Klein J, Holler T, Löffelholz K. Synergistic effect of nicotinamide and choline administration on extracellular choline levels in the brain. *J Pharmacol Exp Ther.* 1993; 266: 720–725.
- (17) Vargas HM, Jenden DJ. Elevation of cerebrospinal fluid choline levels by nicotinamide involves the enzymatic formation of N¹-methylnicotinamide in brain tissue. *Life Sci.* 1996; 58: 1995–2002.
- (18) Orth M, Schapira AHV. Mitochondria and degenerative disorders. *Am J Med Genet.* 2001; 106: 27–36.
- (19) Schulz JB, Matthews RT, Klockgether T, Dichgans J, Beal MF. The role of mitochondrial dysfunction and neuronal nitric oxide in animal models of neurodegenerative diseases. *Mol Cell Biochem.* 1997; 174: 193–197.
- (20) Kosel S, Hofhaus G, Maassen A, Vieregge P, Graeber MB. Role of mitochondria in Parkinson disease. *Biol Chem.* 1999; 380: 865–870.
- (21) Tanaka M, Kovalenko SA, Gong JS, Borgeld HJ, Katsumata K, Hayakawa M, Yoneda M, Ozawa T. Accumulation of deletions and point mutations in mitochondrial genome in degenerative diseases. *Ann N Y Acad Sci.* 1996; 786: 102–111.
- (22) Parker Jr WD, Boyson SJ, Parks JK. Abnormalities of the electron transport chain in idiopathic Parkinson's disease. *Ann Neurol.* 1989; 26: 719–723.

- (23) Schapira AH, Mann VM, Cooper JM, Dexter D, Daniel SE, Jenner P, Clark JB, Marsden CD. Anatomic and disease specificity of NADH CoQ₁ reductase (complex I) in Parkinson's disease. *J Neurochem*. 1990; 55: 2142–2145.
- (24) Swerdlow RH, Parks JK, Miller SW, Tuttle JB, Trimmer PA, Sheehan JP, Bennett JP Jr, Davis RE, Parker WD Jr. Origin and functional consequences of the complex I defect in Parkinson's disease. *Ann Neurol*. 1996; 40: 663–671.
- (25) Mizuno Y, Ohta S, Tanaka M, Takamiya S, Suzuki K, Sato T, Oya H, Ozawa T, Kagawa Y. Deficiencies in complex I subunits of the respiratory chain in Parkinson's disease. *Biochem Biophys Res Commun*. 1989; 163: 1450–1455.
- (26) van der Walt JM, Nicodemus KK, Martin ER, Scott WK, Nance MA, Watts RL, Hubble JP, Haines JL, Koller WC, Lyons K, Pahwa R, Stern MB, Colcher A, Hiner BC, Jankovic J, Ondo WG, Allen FH Jr, Goetz CG, Small GW, Mastaglia F, Stajich JM, McLaurin AC, Middleton LT, Scott BL, Schmechel DE, Pericak-Vance MA, Vance JM. Mitochondrial polymorphisms significantly reduce the risk of Parkinson disease. *Am J Hum Genet*. 2003; 72: 804–811.
- (27) DiDonato S, Zeviani M, Giovannini P, Savarese N, Rimoldi M, Mariotti C, Girotti F, Caraceni T. Respiratory chain and mitochondrial DNA in muscle and brain in Parkinson's disease patients. *Neurology*. 1993; 43: 2262–2268.
- (28) Sandy MS, Langston JW, Smith MT, Di Monte DA. PCR analysis of platelet mtDNA: lack of specific changes in Parkinson's disease. *Mov Disord*. 1993; 8: 74–82.
- (29) Zhang J, Pieper A, Snyder SH. Poly(ADP-ribose) synthetase activation: an early indicator of neurotoxic DNA damage. *J Neurochem*. 1995; 65: 1411–1414.
- (30) Pieper AA, Verma A, Zhang J, Snyder SH. Poly (ADP-ribose) polymerase, nitric oxide and cell death. *Trends Pharmacol Sci*. 1999; 20: 171–181.
- (31) Boulu RG, Mesenge C, Charriaut-Marlangue C, Verrecchia C, Plotkine M. Neuronal death: potential role of the nuclear enzyme, poly (ADP-ribose) polymerase. *Bull Acad Natl Med*. 2001; 185: 555–563.
- (32) Brune B, Dimmeler S, Molina Y, Vedia L, Lapetina EG. Nitric oxide: a signal for ADP-ribosylation of proteins. *Life Sci*. 1994; 54: 61–70.
- (33) Dimmeler S, Brune B. Characterization of a nitric-oxide-catalysed ADP-ribosylation of glyceraldehyde-3-phosphate dehydrogenase. *Eur J Biochem*. 1992; 210: 305–310.
- (34) Zhang J, Snyder SH. Nitric oxide stimulates auto-ADP-ribosylation of glyceraldehyde-3-phosphate dehydrogenase. *Proc Natl Acad Sci U S A*. 1992; 89: 9382–9385.
- (35) Lo EH, Bosque-Hamilton P, Meng W. Inhibition of poly(ADP-ribose) polymerase: reduction of ischemic injury and attenuation of N-methyl-D-aspartate-induced neurotransmitter dysregulation. *Stroke*. 1998; 29: 830–836.
- (36) Takahashi K, Pieper AA, Croul SE, Zhang J, Snyder SH, Greenberg JH. Post-treatment with an inhibitor of poly(ADP-ribose) polymerase attenuates cerebral damage in focal ischemia. *Brain Res*. 1999; 829: 46–54.
- (37) Hageman GJ, Stierum RH, van Herwijnen MH, van der Veer MS, Kleinjans JC. Nicotinic acid supplementation: effects on niacin status, cytogenetic damage, and poly(ADP-ribosylation) in lymphocytes of smokers. *Nutr Cancer*. 1998; 32: 113–120.
- (38) Mandir AS, Przedborski S, Jackson-Lewis V, Wang ZQ, Simbulan-Rosenthal CM, Smulson ME, Hoffman BE, Guastella DB, Dawson VL, Dawson TM. Poly(ADP-ribose) polymerase activation mediates 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced parkinsonism. *Proc Natl Acad Sci U S A*. 1999; 96: 5774–5779.
- (39) Cosi C, Colpaert F, Koek W, Degryse A, Marien M. Poly(ADP-ribose) polymerase inhibitors protect against MPTP-induced depletions of striatal dopamine and cortical noradrenaline in C57B1/6 mice. *Brain Res*. 1996; 729: 264–269.
- (40) Cosi C, Marien M. Decreases in mouse brain NAD⁺ and ATP induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP): prevention by the poly(ADP-ribose) polymerase inhibitor, benzamide. *Brain Res*. 1998; 809: 58–67.
- (41) Shibata K. Nutritional factors affecting the activity of liver nicotinamide methyltransferase and urinary excretion of N¹-methylnicotinamide in rats. *Agric Biol Chem*. 1986; 50: 1489–1493.
- (42) Sano A, Endo N, Takitani S. Fluorometric assay of rat tissue N-methyltransferases with nicotinamide and four isomeric methylnicotinamides. *Chem Pharm Bull*. 1992; 40: 153–156.
- (43) Seifert R, Hoshino J, Kröger H. Nicotinamide methylation: tissue distribution, development and neoplastic changes. *Biochim Biophys Acta*. 1984; 801: 259–264.
- (44) Fukushima T, Kaetsu A, Lim H, Moriyama M. Possible role of 1-methylnicotinamide in the pathogenesis of Parkinson's disease. *Exp Toxicol Pathol*. 2002; 53: 469–473.
- (45) Matsubara K, Aoyama K, Suno M, Awaya T. N-methylation underlying Parkinson's disease. *Neurotoxicol Teratol*. 2002; 24: 593–598.
- (46) Parsons RB, Smith SW, Waring RH, Williams AC, Ramsden DB. High expression of nicotinamide N-methyltransferase in patients with idiopathic Parkinson's disease. *Neurosci Lett*. 2003; 342: 13–16.
- (47) Willets JM, Lunec J, Williams AC, Griffiths HR. Neurotoxicity of nicotinamide derivatives; their role in the aetiology of Parkinson's disease. *Biochem Soc Trans*. 1993; 21: 299S.
- (48) Fukushima T, Tawara T, Isobe A, Hojo N, Shiwaku K, Yamane Y. Radical formation site of cerebral complex I and Parkinson's disease. *J Neurosci Res*. 1995; 42: 385–390.
- (49) Narabayashi H, Takeshige K, Minakami S. Alteration of inner-membrane components and damage to electron-transfer activities of bovine heart submitochondrial particles induced by NADPH-dependent lipid peroxidation. *Biochem J*. 1982; 202: 97–105.
- (50) Chomyn A, Mariottini P, Cleeter MW, Ragan CI, Matsuno-Yagi A, Hatefi Y, Doolittle RF, Attardi G. Six unidentified reading frames of human mitochondrial DNA encode components of the respiratory-chain NADH dehydrogenase. *Nature*. 1985; 314: 592–597.
- (51) Triepels RH, Van Den Heuvel LP, Trijbels JM, Smeitink JA. Respiratory chain complex I deficiency. *Am J Med Genet*. 2001; 106: 37–45.
- (52) Peter AM. Structures and function of the water-soluble vitamins. In: David AB editor. *Harper's Biochemistry 25th edition*, Stamford: Appleton & Lange, 2000: p. 676–678.
- (53) Food and Agriculture Organization of the United Nations. Rome Declaration on World Food Security and World Food Summit Plan of Action. Technical background documents 1–5. 1996.
- (54) Cosnett JE, Bill PL. Parkinson's disease in blacks. Observa-

- tions on epidemiology in Natal. *S Afr Med J.* 1988; 73: 281–283.
- (55) Lombard A, Gelfand M. Parkinson's disease in the African. *Cent Afr J Med.* 1978; 24: 5–8.
- (56) Kusumi M, Nakashima K, Harada H, Nakayama H, Takahashi K. Epidemiology of Parkinson's disease in Yonago City, Japan: comparison with a study carried out 12 years ago. *Neuroepidemiology.* 1996; 15: 201–207.
- (57) Fukushima T, Moriyama M. Corn might prevent Parkinson's disease. *Clin Nutr.* 2001; 20: 559.
- (58) Fukushima T, Tanaka K, Ushijima K, Moriyama M. Retrospective study of preventive effect of maize on mortality from Parkinson's disease in Japan. *Asia Pac J Clin Nutr.* 2003; 12: 447–450.
- (59) National Nutrition Survey, Japan. The Ministry of Health and Welfare, Japan.